



Individual whey protein components influence lipid oxidation dependent on pH

Horn, Anna Frisenfeldt; Nielsen, Nina Skall; Jacobsen, Charlotte

Publication date:
2012

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Horn, A. F., Nielsen, N. S., & Jacobsen, C. (2012). *Individual whey protein components influence lipid oxidation dependent on pH*. Poster session presented at 103rd AOCS Annual Meeting & Expo, Long Beach, CA, United States.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Conference abstract (Poster presentation)

103rd AOCS Annual Meeting & Expo, April 29-May 2, 2012, Long Beach, California, USA

Individual whey protein components influence lipid oxidation dependent on pH

Anna Frisenfeldt Horn, Nina Skall Nielsen & Charlotte Jacobsen

National Food Institute, Technical University of Denmark

In emulsions, lipid oxidation is expected to be initiated at the oil-water interface. The properties of the emulsifier used and the composition at the interface is therefore expected to be of great importance for the resulting oxidation. Previous studies have shown that individual whey protein components (α -lactalbumin and β -lactoglobulin) adsorb differently to the interface depending on pH. In addition, differences has been shown to exists between the oxidative stability provided by α -lactalbumin and β -lactoglobulin. The hypothesis is that pH influences the oxidative stability of emulsions by affecting the preferential adsorption of whey protein components at the interface.

The aim of the study was to compare lipid oxidation in 10% fish oil-in-water emulsions prepared with 1% whey protein having either a high concentration of α -lactalbumin, a high concentration of β -lactoglobulin or equal amounts of the two. Emulsions were prepared at pH4 and pH7. Emulsions were characterized by their droplet sizes, viscosities, and contents of proteins in the water phase. Lipid oxidation was assessed by PV and secondary volatile oxidation products. Results showed that pH greatly influenced the oxidative stability of emulsions. At high pH β -lactoglobulin emulsions were more stable than α -lactalbumin emulsions, whereas at low pH the opposite was the case.